

Investigation of Dust-Ion Acoustic Waves in a Magnetized Collisional Dusty Plasma with Kappa Distribution Function for Electrons

Hossein Zahed^{a,*}, Mohammad Reza Sayyar^{b,c}, Sayyed Jalal Pestehe^b, and Samad Sobhanian^b

^aDepartment of Physics, Sahand University of Technology, Tabriz, Iran

^bDepartment of Atomic and Molecular Physics, University of Tabriz, Tabriz, Iran

^cFaculty of Science, University of Maragheh, Maragheh, Iran

*Corresponding author' Email: Zahed@sut.ac.ir

Received: Aug. 16, 2016, Revised: Jan. 29, 2017, Accepted: Apr. 26, 2017, Available Online: Dec. 1, 2018

DOI: 10.29252/ijop.12.2.81

ABSTRACT— The propagation of arbitrary amplitude dust ion acoustic waves (DIAWs) in a magnetized collisional dusty plasma including hot electrons, with kappa velocity distribution for electrons, warm ions and dust particles has been studied. In the presence of immobile massive dust particulates, DIAWs have been investigated through the Sagdeev pseudo-potential method. It is demonstrated that the amplitude and width of the pseudo-potential are increased with the ion density and also with Directional cosines. It is shown that the behaviors of the amplitude and the width of the wave in terms of all of plasma parameters is similar to the our recently work, and the spectral index has a little effect on the wave.

KEYWORDS: magnetized dusty plasma, Dust ion acoustic waves, Sagdeev pseudo-potential method.

I. INTRODUCTION

Nowadays, the study of dusty plasmas properties has received a great deal of attention both theoretically and experimentally. In the presence of massive and highly charged dust particulates in an usual electron ion plasma new types of waves, such as dust acoustic (DA) waves and dust-ion-acoustic (DIA) waves are excited. These dusts can be regarded as static or mobile particles [1]. The nonlinear waves particularly, the DIA solitary waves (DIASW) have been theoretically investigated by several authors [2-6].

In the most of the above mentioned investigations, Maxwellian or Maxwellian Boltzmann distribution functions have been used. However, a lot of theoretical observations of space plasmas are often characterized by a non-Maxwellian particle distribution function. In these plasmas, superthermal particles which are produced due to the effect of wave particle interaction or external forces. Superthermal plasmas are relativistic pulsar wind, solar wind, magnetosphere, interstellar medium, auroral zone plasmas, plasmas produced during an ultra-intense laser pulse interaction with matter. [7, 8]. These kinds of a plasma, can be characterized by generalized Lorentzian or kappa distribution function [9, 10]. This Lorentzian (kappa) velocity distribution is used to model the electrons in magnetosphere and electromagnetic ion cyclotron waves in equatorial ring protons [11, 12]. Using a kappa distribution for plasma particles, many authors have studied the propagation of ion acoustic waves in a magnetized plasma using Sagdeev potential method. The one dimensional kappa velocity distribution is

$$F_{\kappa}(v) = \frac{1}{(\pi\kappa\theta^2)^{3/2}} \frac{\Gamma(\kappa+1)}{\Gamma(\kappa-1/2)} \left(1 + \frac{v^2}{\kappa\theta^2}\right)^{-(\kappa+1)} \quad (1)$$

where θ is the most probable speed (effective thermal speed), related to the usual thermal

velocity $V_i = (K_B T / m)^{1/2}$ by $\theta = [(2\kappa - 3)/\kappa]V_i$, T is the characteristic kinetic temperature, κ is spectral index and K_B is the Boltzmann constant. The most probable speed, and hence the κ -distribution, is defined for $\kappa > 3/2$ [7, 8, 11]. Nonlinear propagation of ion acoustic waves in a magnetized plasma, electron acoustic waves in a two temperature electron plasma and linear and nonlinear propagation of these sound waves again with the presence of hot electron component and so on have been already studied [13].

Recently, we have investigated the propagation of dust ion acoustic wave in a dusty plasma in the presence of nonthermal electrons [14]. In this paper, the previous work has been extended with the warm ions and hot electrons. In the second section, the governing basic equations are presented similar to our previous paper [14]. The resultant dispersion relation and as well as the Sagdeev potential are derived in the third section. Finally, the numerical investigations are presented in the last section.

II. BASIC EQUATIONS

The continuity and momentum equations for ions and also Poisson equation are:

$$\partial_t n_i + \nabla \cdot (n_i \mathbf{v}_i) = 0, \quad (2)$$

$$\partial_t \mathbf{v}_i + (\mathbf{v}_i \cdot \nabla) \mathbf{v}_i = \frac{e}{m_i} \left(\mathbf{E} + \frac{B_0 \mathbf{v}_i \times \mathbf{e}_z}{c} \right) - \nabla \mathbf{p}_i - m_i n_i \mathbf{v}_i \nu, \quad (3)$$

$$\nabla^2 \varphi = -4\pi [-en_e + en_i - ez_d n_d], \quad (4)$$

where n_e , n_i , and n_d are electron, ion, and dust densities respectively. \mathbf{v}_i , ν , and m_i indicating the ion velocity, collision frequency and its mass. $qd = -ezd$ where φ is the plasma potential, z_d is the dust charge number, so that the charge of the dust is given by $qd = -ezd$, with e being the elementary charge. Here, electrons are assumed to be kappa distributed and the expression for the electron density is as follows:

$$n_e = \frac{1}{\delta_1} \left(1 - \frac{\varphi}{\kappa - 3/2} \right)^{-(\kappa - 1/2)}, \quad (5)$$

$$\text{where, } \delta_1 = \frac{n_{i0}}{n_{e0}}.$$

We assume that the wave is propagating in the x - z plane. After normalization, the above system of equations is reduced to:

$$\partial_t n + \partial_x (n v_x) + \partial_z (n v_z) = 0 \quad (6)$$

$$\partial_t v_x + (v_x \partial_x + v_z \partial_z) v_x = -\partial_x \varphi + v_y - \gamma \frac{\partial_x n}{n} + \nu v_x \quad (7)$$

$$\partial_t v_y + (v_x \partial_x + v_z \partial_z) v_y = -v_x - \gamma \frac{\partial_y n}{n} + \nu v_y \quad (8)$$

$$\partial_t v_z + (v_x \partial_x + v_z \partial_z) v_z = -\partial_z \varphi - \gamma \frac{\partial_z n}{n} + \nu v_z \quad (9)$$

$$(\partial_{xx} + \partial_{zz}) \varphi = \beta \left[\frac{1}{\delta_1} \left(1 - \frac{\varphi}{\kappa - 3/2} \right)^{-(\kappa - 1/2)} - \delta_1 n + \delta_2 \right] \quad (10)$$

where, $\beta = r_g^2 / \lambda_e^2$, $\delta_1 = n_{i0} / n_{e0}$, $\delta_2 = n_d z_d / n_{e0}$. $r_g = C_s / \Omega$ is the ion gyro radius and $\lambda_e = (T_e / 4\pi n_{e0} e^2)^{1/2}$ is the electron Debye length. $C_s = (T_e / m_i)^{1/2}$ is the ion acoustic velocity, and $\Omega = eB_0 / m_i c$ is the ion gyro frequency. n_{e0} and n_{i0} are unperturbed electron and ion densities, respectively. The normalizations are:

$$\Omega t \rightarrow t, (C_s / \Omega) \nabla \rightarrow \nabla, v_i / C_s \rightarrow v,$$

$$n_i / n_{i0} \rightarrow n, \text{ and } e\varphi / T_e \rightarrow \varphi,$$

To derive the dispersion relation for low frequency waves, we write the dependent variables as sum of equilibrium and perturbed parts. Assuming $n = 1 + \bar{n}$, $v_x = \bar{v}_x$, $v_z = \bar{v}_z$, $v_y = \bar{v}_y$, Eqs. (5) - (9) can be rewritten as:

$$\partial_t \bar{n} + \partial_x \bar{v}_x + \partial_z \bar{v}_z = 0 \quad (11)$$

$$\partial_t \bar{v}_x = -\partial_x \bar{\phi} + \bar{v}_y - \gamma \frac{\partial_x \bar{n}}{1+n} + \nu v_x \quad (12)$$

$$\partial_t \bar{v}_y = -v_x + \nu v_y \quad (13)$$

$$\partial_t \bar{v}_z = -\partial_z \bar{\phi} - \gamma \frac{\partial_z \bar{n}}{1+n} + \nu v_z \quad (14)$$

$$(\partial_{xx} + \partial_{zz}) \bar{\phi} = \beta \left[\frac{\kappa - 1/2}{\kappa - 3/2} \bar{\phi} - \delta_1 \bar{n} \right] \quad (15)$$

We assume that the perturbation is of the form $\exp i(k_x x + k_z z - \omega t)$, where k_x and k_z are the wave numbers in x and z directions, respectively. Hence, the dispersion relation for ion acoustic wave can be obtained as

$$\beta_4 (i\omega(\nu + i\omega)^3 + k^2(\nu + i\omega)^2 \gamma + i\omega(\nu + i\omega) + \gamma k_z^2 - \delta_1 k^2(\nu + i\omega)^2 - \delta_1 k_z^2) = 0 \quad (16)$$

$$\begin{aligned} & (\gamma \nu^2 k^2 - (\gamma \omega^2 - \gamma) k_z^2 - 3\nu^2 \omega^2 + \omega^4 - \omega^2) \\ & - \delta_1 \left((\nu^2 + \omega^2) k^2 - k_z^2 / \beta_4 + i(2\gamma \nu \omega k^2 \right. \\ & \left. + \nu^3 \omega - 3\nu \omega^3 + \nu \omega) - 2\delta_1 \nu \omega k^2 \right) = 0 \end{aligned} \quad (17)$$

where $\beta_4 = k^2 / \beta + \frac{\kappa - 1/2}{\kappa - 3/2}$. By assuming $\omega = \omega_r + i\omega_i$, the above equation reads as

$X + iY = 0$, where

$$\begin{aligned} X = \beta_4 \left((1 + 3\nu^2) \lambda + \gamma k_z^2 + \gamma k^2 (\nu^2 + \lambda) \right. \\ \left. - \omega_i \nu (\nu^2 + 1 - 6\omega_r^2 + 3\lambda + 2\gamma k^2) \right) \\ + \delta_1 k^2 (\nu^2 - 2\nu \omega_i \lambda) - \delta_1 k_z^2 \end{aligned} \quad (18)$$

and

$$\begin{aligned} Y = -2\omega_r \omega_i k^2 (1 + 3\nu^2) (\gamma + \delta_1) \\ + \beta_4 [\nu^3 \omega_r - 3\nu \omega_r^3 + 9\nu \omega_r \omega_i^2 \\ + \nu \omega_r (1 + 2\gamma k^2) + 2\delta_1 \nu \omega_r k^2] \end{aligned} \quad (19)$$

Here $\lambda = \omega_i^2 - \omega_r^2$ introducing a new variable $\eta = l_x x + l_z z - Mt$, where l_x and l_z are directional cosines and M is the Mach number. Equations (5)–(9) can be written as ordinary differential equations in terms of η :

$$M(1 - n) + n(l_x v_x + l_z v_z) = 0 \quad (20)$$

$$\begin{aligned} -M \frac{dv_x}{d\eta} + \left(v_x l_x \frac{d}{d\eta} + v_z l_z \frac{d}{d\eta} \right) v_x \\ = -l_x \frac{d\phi}{d\eta} + v_y - \gamma l_x \frac{dn}{nd\eta} + \nu v_x \end{aligned} \quad (21)$$

$$\begin{aligned} -M \frac{dv_y}{d\eta} + \left(v_x l_x \frac{d}{d\eta} + v_z l_z \frac{d}{d\eta} \right) v_y \\ = -v_x \gamma \frac{dn}{nd\eta} + \nu v_y \end{aligned} \quad (22)$$

$$\begin{aligned} -M \frac{dv_y}{d\eta} + \left(v_x l_x \frac{d}{d\eta} + v_z l_z \frac{d}{d\eta} \right) v_y \\ = -v_x \gamma \frac{dn}{nd\eta} + \nu v_y \end{aligned} \quad (23)$$

$$\begin{aligned} -M \frac{dv_z}{d\eta} + \left(v_x l_x \frac{d}{d\eta} + v_z l_z \frac{d}{d\eta} \right) v_z \\ = -l_z \frac{d\phi}{d\eta} - \gamma l_z \frac{dn}{nd\eta} + \nu v_z \end{aligned} \quad (24)$$

Substituting Eq. (19) in Eq. (20), we have:

$$\frac{M}{n} \frac{dv_x}{d\eta} = l_x \frac{d\phi}{d\eta} - v_y + \gamma l_x \frac{dn}{nd\eta} + \nu v_x \quad (25)$$

Using Eq. (19), Eqs. (21) and (20) are rewritten as:

$$\frac{M}{n} \frac{dv_y}{d\eta} = v_x + \nu v_y \quad (26)$$

$$\frac{M}{n} \frac{dv_z}{d\eta} = l_z \frac{d\phi}{d\eta} + \gamma l_z \frac{dn}{nd\eta} + \nu v_z \quad (27)$$

$$G(\phi) = \int_0^\phi nd\phi \quad (28)$$

$$v_z = b_1\varphi + b_2\varphi^2 + b_3\varphi^3 + \dots \quad (29)$$

Considering Eq. (23), we get:

$$\frac{1}{2}(l_x^2 + l_z^2) \left(\frac{d\varphi}{d\eta}\right)^2 = \beta \left[1 + \varphi + \frac{\kappa - 1/2}{2(\kappa - 3/2)} \varphi^2 - \delta_1 G(\varphi) + \delta_2 \varphi \right] \quad (10)$$

Comparing the coefficients of equal powers of φ in Eqs. (26) and (28), up to the second order, one can obtain:

$$\left(\frac{1}{3} \frac{(-2\gamma a_1 l_z + M b_1 - l_z) A_2}{\sqrt{A_1}} + (-6\gamma a_2 l_z + 2M b_2 - 2a_1 l_z) \sqrt{A_1} - 2a_1 v b_1 - v b_2 \right) \varphi^2 + ((-2\gamma a_1 l_z + M b_1 - l_z) + \sqrt{A_1} - v b_1) \varphi = 0 \quad (21)$$

III. PSEUDO-POTENTIAL APPROACH

The Eq. (29) can be written as

$$\frac{1}{2} \left(\frac{d\varphi}{d\eta}\right)^2 + V(\varphi) = 0 \quad (32)$$

The above equation is an “energy integral” of an oscillatory particle with unit mass, pseudo-velocity $\frac{d\varphi}{d\eta}$ at the pseudo-position φ in a pseudo-potential well $V(\varphi)$. To obtain an equation for $G(\varphi)$, one can expand $G(\varphi)$ and v_x in terms of φ :

$$G(\varphi) = \varphi + a_1\varphi^2 + a_2\varphi^3 + \dots \quad (33)$$

$$v_x = A\varphi + B\varphi^2 + C\varphi^3 + \dots \quad (34)$$

To find the coefficients, one can substitute the Eqs.(31) and (32) in Eq. (29) to get,

$$\left(\frac{d\varphi}{d\eta}\right)^2 = \frac{2\beta}{l_x^2 + l_z^2} \left[(1 + \delta_1 - \delta_2)\varphi + \left(\frac{2(\kappa - 1/2)}{\kappa - 3/2} + 4\delta_1 a_1\right)\varphi^2 + \left(\frac{\kappa^2 - 1/4}{(\kappa - 3/2)^2} - 6\delta_1 a_2\right)\varphi^3 + \dots \right], \quad (45)$$

And up to the second order we have

$$\frac{d^2\varphi}{d\eta^2} = A_1\varphi + A_2\varphi^2 = -\frac{dV}{d\varphi} \quad (56)$$

$$\frac{d^2\varphi}{d\eta^2} = A_1\varphi + A_2\varphi^2 = -\frac{dV}{d\varphi} \quad (67)$$

where,

$$A_1 = \frac{\beta}{l_x^2 + l_z^2} \left[\frac{2(\kappa - 1/2)}{\kappa - 3/2} + 4\delta_1 a_1 \right] \quad (38)$$

and

$$A_2 = \frac{\beta}{l_x^2 + l_z^2} \left[\frac{\kappa^2 - 1/4}{(\kappa - 3/2)^2} - 6\delta_1 a_2 \right] \quad (79)$$

Eqs. (24) and (25) can be solved as

$$M \frac{dv_x}{d\varphi} \frac{d\varphi}{d\eta} = v_x G'(\varphi) + v v_y G'(\varphi) \quad (40)$$

$$v_y = l_x \frac{d\varphi}{d\eta} + l_x \frac{\gamma}{G'(\varphi)} \frac{dG'(\varphi)}{d\eta} + v v_x - \frac{M}{G'(\varphi)} \frac{dv_x}{d\eta} \quad (41)$$

Equating the coefficients in the Eqs. (36) and (37), and expansion of $\left(\frac{d\varphi}{d\eta}\right)$ and substituting in Eq. (39) one can obtains up to second order.

$$\begin{aligned}
 & \left(\frac{10l_z a_1 b_1}{l_x} - \frac{16M v^2 a_1^2}{l_x} - \frac{3M v^2 a_2}{l_x} + \frac{v^2 b_2 l_z}{l_x} \right. \\
 & + 8M a_1 l_x A_1 + \frac{8M^3 A_1 a_1^2}{l_x} - \frac{12M^3 A_1 a_2}{l_x} - \\
 & \frac{2M^3 A_2 a_1}{l_x} - \frac{2l_z b_2 \sqrt{A_1}}{l_x} - 10v a_1 l_x \sqrt{A_1} - \\
 & \frac{1}{3} \frac{v l_x A_2}{\sqrt{A_1}} + \frac{M^2 l_z A_2 b_1}{l_x} + \frac{4M^2 l_z A_1 b_2}{l_x} + \frac{10v^2 a_1 b_1 l_z}{l_x} - \\
 & 12M \gamma a_2 l_x A_1 - 2M \gamma a_1 l_x A_2 - \frac{1}{3} \frac{b_1 l_z A_2}{l_x \sqrt{A_1}} + \\
 & \frac{16M^2 v a_1^2 \sqrt{A_1}}{l_x} + \frac{12M^2 v a_2 \sqrt{A_1}}{l_x} - \frac{4l_z a_1 b_1 \sqrt{A_1}}{l_x} + \\
 & 16\gamma v a_1^2 l_x \sqrt{A_1} + 6\gamma v a_2 l_x \sqrt{A_1} - 12M \gamma a_1^2 l_x A_1 + \\
 & \frac{4M^2 l_z a_1 b_1 A_1}{l_x} + \frac{4}{3} \frac{M^2 v a_1 A_2}{l_x \sqrt{A_1}} + \frac{2}{3} \frac{\gamma v a_1 l_x A_2}{\sqrt{A_1}} + \\
 & \left. \frac{l_z b_2}{l_x} - \frac{16M a_1^2}{l_x} + M l_x A_2 - \frac{3M a_2}{l_x} \right) \varphi^2 + \\
 & \left(-\frac{2M a_1}{l_x} - \frac{2M v^2 a_1}{l_x} + 2\gamma v a_1 l_x \sqrt{A_1} - \frac{2M^3 a_1 A_1}{l_x} + \right. \\
 & \frac{4M^2 v a_1 \sqrt{A_1}}{l_x} + \frac{M^2 l_z b_1 A_1}{l_x} + M l_x A_1 - 2M \gamma a_1 l_x A_1 + \\
 & \left. \frac{v^2 b_1 l_z}{l_x} - \frac{b_1 l_z \sqrt{A_1}}{l_x} + \frac{b_1 l_z}{l_x} - v l_x \sqrt{A_1} \right) \varphi = 0
 \end{aligned} \tag{42}$$

The coefficients a_1 , a_2 , b_1 and b_2 can be determined by solving the Eqs. (30) and (39). From equation (34), as like as [14], we get:

$$\varphi = -\frac{3A_1}{2A_2} \operatorname{sech}^2 \left(\sqrt{A_1} \eta / 2 \right) \tag{43}$$

IV. NUMERICAL INVESTIGATIONS

The above results i.e. Eqs. (35)–(41), show that all coefficients are dependent to the spectral index and all plasma parameters. So for investigating the plasma parameters effects on the wave propagation, we use parameters written in Table 1:

$$\beta = \frac{r_g^2}{\lambda_e^2}, \quad \delta_1 = \frac{n_{i0}}{n_{e0}}$$

TABLE 1: PLASMA PARAMETERS USED IN NUMERICAL INVESTIGATION.

Plasma parameters	β	δ_1	Mach number (M)	Directional cosines (l_z)	Spectral index (κ)
Range of Values	0.46, 0.49, 0.55	1.54, 2.54, 3.54	2, 2.1, 2.25	0.1, 0.4, 0.58	5/2

The variation of the pseudo-potential versus ion to electron density ratio is shown in Fig. 1. It is seen that the amplitude of $V(\varphi)$ increases with increasing the density ratio. It is concluded that increasing the ion density can lead to higher amplitude waves and also it can be said that for all positive values of φ , $V(\varphi)$ is positive and vice versa.




Fig. 1 Variation of Sagdeev potential with plasma potential for different values of δ_1 and for $l_z = 0.1$, $M = 2.25$, and $\beta = 0.46$.

From Fig. 2, it is concluded that the behavior of the pseudo-potential is the same for all β values and only their amplitudes increases slightly. In other words, it can be said that the behavior of the potential is linearly dependent to the electron temperature.

In Fig. 3, the pseudo-potential is plotted as a function of φ . It is clear that for fixed value of the φ , the amplitude of the Sagdeev potential is increasing with l_z .




Fig. 2 The plot of $V(\varphi)$ vs. φ , for $l_z = 0.1$, $M = 2.25$, and $\delta_1 = 2.54$ for different values of β .




Fig. 3 The plot of $V(\varphi)$ vs. φ , for $M = 2.25$, $\delta_1 = 2.54$, and $\beta = 0.46$ for different values of l_z .

Figure 4 shows the plot of pseudo-potential amplitude versus plasma potential for different values of Mach number.




Fig. 4 The plot of $V(\varphi)$ vs. φ , for $l_z = 0.1$, $\delta_1 = 2.54$, and $\beta = 0.46$ for different values of M .

In Fig. 5 the pseudo-potential, φ is plotted versus the η , for $l_z = 0.1$, $M = 2.25$, and $\beta = 0.46$ for different values of δ_1 . It is concluded that increasing the δ_1 may increase the plasma amplitude. Also this is negative for negative values of φ , and vice versa. Figure 6 shows that for all values of δ_1 , i.e. the ratio of ion density to the equilibrium electron density, the plasma potential is negative. It is also seen that the amplitude and the width of the potential increases as δ_1 increases. Comparing the results for high magnitudes of δ_1 , it can be said that the slope of the amplitude is decreased with increasing the ion density.




Fig. 5 The variation of φ vs. η , for $l_z = 0.1$, $M = 2.25$, and $\beta = 0.46$ for different values of δ_1 .




Fig. 6 The plot φ vs. η , for $l_z = 0.1$, $M = 2.25$, and $\delta_1 = 2.54$ different values of β .

The variation of plasma potential is plotted as a function of η for $l_z = 0.1$, $M = 2.25$, and $\delta_1 = 2.54$ for different values of β in Fig. 6. It is seen that decreasing the magnitude of β may increase the width and amplitude of the potential.

In Fig. 7, we have illustrated the variation of plasma potential as a function of η for $M = 2.25$, $\delta_1 = 2.54$, and $\beta = 0.46$ for different values of l_z and for values of δ_1 . It is seen that the amplitude is negative and that the negative amplitude and the width of the potential are increased by increasing l_z .




Fig. 7 The plot of ϕ vs. η , for $M = 2.25$, $\delta_1 = 2.54$, $\beta = 0.46$ for different values of l_z .




Fig. 8 The plot of ϕ vs. η , for $l_z = 0.1$, $\delta_1 = 2.54$, and $\beta = 0.46$ for different values of M .

In Fig. 8, we have illustrated the variation of plasma potential as a function of η for $l_z = 0.1$,

$\delta_1 = 2.54$, and $\beta = 0.46$ for different values of Mach number.

V. CONCLUSION

Using Sagdeev potential, the dispersion relation of dust ion acoustic wave in a collisional magnetized dusty plasma is obtained in the presence of warm ions and hot electrons. It is shown that the amplitude and the width of the wave is dependent on the plasma parameters such as plasma particles density and temperatures, plasma potential, Mach number and etc. The behavior of the amplitude and the width of the wave in terms of all of these parameters is similar to that in our previous work [14]. However, the values of these two variables are different. It is shown that the Sagdeev potential has negative value for positive plasma potential and vice versa. We have also shown that increasing δ_1 and decreasing β can lead to the increasing of wave amplitude and also they can cause an increase in the width and the amplitude of the potential. Finally, it is found that the spectral index, i.e. the quantity when goes to infinite, the kappa distribution changes to Maxwellian one, has little effect on the propagation properties of the wave.

REFERENCES

- [1] M. Alam, M. Masud, and A. Mamun, "Effects of two-temperature superthermal electrons on dust-ion-acoustic solitary waves and double layers in dusty plasmas," *Astrophysics Space Sc.*, vol. 349, pp. 245-253, 2014.
- [2] S. El-Labany, E. El-Shamy, and S. El-Warraki, "Effects of two-temperature electrons, external oblique magnetic field, and higher-order nonlinearity on dust acoustic solitary waves in a dusty plasma with vortex-like ion distribution," *Astrophysics Space Sc.*, vol. 315, pp. 287-295, 2008.
- [3] M. Hossain, A. Mamun, and K. Ashrafi, "Cylindrical and spherical dust ion-acoustic Gardner solitons in a quantum plasma," *Phys. Plasmas*, vol. 18, pp. 103704 (1-6), 2011.
- [4] S. Popel, A. Golub, T. Losseva, A. Ivlev, S. Khrapak, and G. Morfill, "Weakly dissipative dust-ion-acoustic solitons," *Phys. Rev. E*, vol. 67, pp. 056402 (1-5), 2003.

- [5] N. Kundu, M. Masud, K. Ashrafi, and A. Mamun, "Dust-ion-acoustic solitary waves and their multi-dimensional instability in a magnetized nonthermal dusty electronegative plasma," *Astrophys. Space Sci.* vol. 343, pp. 279-287, 2013.
- [6] A. Mamun, N. Jahan, and P. Shukla, "DIA and DA solitary waves in adiabatic dusty plasmas," *J. Plasma Phys.* vol. 75, pp. 413-431, 2009.
- [7] T.K. Baluku and M.A. Hellberg, "Ion acoustic solitons in a plasma with two-temperature kappa-distributed electrons," *Phys. Plasmas*, vol. 19, pp. 012106 (1-10), 2012.
- [8] A. Saha and P. Chatterjee, "New analytical solutions for dust acoustic solitary and periodic waves in an unmagnetized dusty plasma with kappa distributed electrons and ions," *Phys. Plasmas*, vol. 21, pp. 02211 (1-6), 2014.
- [9] T.K. Baluku and M.A. Hellberg, "Dust acoustic solitons in plasmas with kappa-distributed electrons and/or ions," *Phys. Plasmas*, vol. 15, pp. 123705 (1-11), 2008.
- [10] M.A. Hellberg, R.L. Mace, T.K. Baluku, I. Kourakis, and N. Saini, "Comment on "Mathematical and physical aspects of Kappa velocity distribution" [*Phys. Plasmas* 14, 110702 (2007)]," *Phys. Plasmas*, vol. 16, pp. 094701 (1-9), 2009.
- [11] G. Banerjee and S. Maitra, "Pseudopotential approach for dust acoustic solitary waves in dusty plasmas with kappa-distributed ions and electrons and dust grains having power law size distribution," *Phys. Plasmas* (1994-present), vol. 22, pp. 043708-9, 2015.
- [12] N. Saini, N. Kaur, and T.S. Gill, "Dust acoustic solitary waves of Kadomstev-Petviashvili (KP) equation in superthermal dusty plasma," *Advances in Space Research*, vol. 55, pp. 2873-2882, 2015.
- [13] M. Michael, S. Gopinathan, S. Sebastian, N. T. Willington, A. Varghese, R. Gangadharan, and Ch. Venugopal "Kadomstev-Petviashvili-Burgers (KPB) Equation in a Five Component Cometary Plasma with Kappa Described Electrons and Ions," *J. Adv. Med. Education Profession.*, vol. 3, pp. 1431-1442, 2015.
- [14] M. R. Sayyar, H. Zahed, S. J. Pestehe, and S. Sobhanian, "Propagation of arbitrary amplitude dust-ion acoustic waves in a

collisional magnetized dusty plasma in the presence of non-thermal electrons", *Phys. Plasmas*, vol. 23, pp. 73704 (1-6), 2016.



Samad Sobhanian, Professor Emeritus in plasma physics, from University of Tabriz, is graduated from University of Paul Sabatier, Toulouse, France in atomic collisions in plasma. His main research works include both theoretical and experimental plasma physics. He is a member of Iran Nanotechnology, Physics society of Iran and Optics and Photonics Society of Iran.



Hossein Zahed, associate professor in plasma physics, from Sahand University of Technology, is graduated from University of Tabriz in plasma. His main research works is theoretical plasma physics. He is a member of Iran Nanotechnology, Physics Society of Iran and Optics and Photonics Society of Iran.



Sayyed Jalal Pestehe is associated professor of atomic and molecular department of the University of Tabriz. He completed his B.Sc. and M.Sc. and Ph.D. studies at the Universities of Tabriz and York, respectively. He has supervised 24 M.Sc. and 3 Ph.D. thesis since 2002 and published nearly 80 papers in journals and conference proceedings. His

recent work focuses on the low temperature plasmas, nanotechnology, and dusty plasmas. He is a member of Iran nanotechnology, Physics Society of Iran and Optics and Photonics Society of Iran.



Mohammad Reza Sayyar, assistant professor of Plasma Physics in physics department of the

Maragheh University, He is graduated from University of Tabriz in plasma physics. He is a member of Iran Nanotechnology, Physics Society of Iran and Optics and Photonics Society of Iran.

THIS PAGE IS INTENTIONALLY LEFT BLANK.